

HECKE ALGEBRAS, MODULAR CATEGORIES AND 3-MANIFOLDS QUANTUM INVARIANTS

CHRISTIAN BLANCHET

ABSTRACT. We construct modular categories from Hecke algebras at roots of unity. For a special choice of the framing parameter, we recover the Reshetikhin-Turaev invariants of closed 3-manifolds constructed from the quantum groups $U_q sl(N)$ by Reshetikhin-Turaev and Turaev-Wenzl, and from skein theory by Yokota. The possibility of such a construction was suggested by Turaev, as a consequence of Schur-Weil duality. We then discuss the choice of the framing parameter. This leads, for any rank N and level K , to a modular category $\tilde{H}^{N,K}$ and a reduced invariant $\tilde{\tau}_{N,K}$. If N and K are coprime, then this invariant coincides with the known invariant $\tau^{PSU(N)}$ at level K . If $\gcd(N, K) = d > 1$, then we show that the reduced invariant admits spin or cohomological refinements, with a nice decomposition formula which extends a theorem of H. Murakami.

INTRODUCTION

Our goal is to construct modular categories from the Hecke algebras of type A at roots of unity, and to obtain certain reductions and refinements of them. Our main results are the followings.

- We give a (reasonably self contained) construction of modular categories underlying the known $SU(N)$ quantum invariants. As expected, at rank-level (N, K) , isomorphism classes of simple objects are indexed by the set of Young diagrams with at most $N - 1$ lines and K columns, whose cardinality is $\frac{(N+K-1)!}{(N-1)!K!}$. The dimensions of the TQFT modules are given by the Verlinde formula.
- At rank-level (N, K) we obtain a reduced modular category $\tilde{H}^{N,K}$, and a reduced invariant $\tilde{\tau}_{N,K}$. Here the number of non-isomorphic simple objects is $\frac{(N+K-1)!}{N!K!}d$, with $d = \gcd(N, K)$. Except for $\gcd(N, K) = 1$, in which case we recover the $PSU(N)$ invariant, this result seems to be new.

Date: February 1998.

Key words and phrases. Hecke algebra, 3-manifold, quantum invariant, TQFT, modular functor, modular category.

- We use a grading modulo $d = \gcd(N, K)$ to define invariants of 3-manifolds equipped with what we call a spin structure with coefficients modulo d , and we prove a decomposition formula which extends a theorem of H. Murakami [29]. This refinement holds when d is even, and $\frac{N}{d}, \frac{K}{d}$ are odd (the spin case); in the other cases, we obtain cohomological refinements.

The quantum invariants, predicted by Witten using Chern-Simons theory and path integrals, were first constructed by Reshetikhin-Turaev [33] and Turaev-Wenzl [36] using representation theory of quantum groups. The work of Turaev [35] shows that a key concept in the construction of these invariants, as well as in extending them to a topological quantum field theory (TQFT), is that of a modular category.

A modular category is a braided category with some additional algebraic features (duality, twist, a finite set of simple objects satisfying a domination property and a non-degeneracy axiom). The interest of this concept is that it provides a topological quantum field theory in dimension 3, and in particular, invariants of links and 3-manifolds. However, in particular examples, it is not easy to define precisely the modular category and to check the required properties. For the $SU(N)$ invariants, first constructed using representation theory of $U_q(sl(N))$, it is known that underlying modular categories can be derived from the category of representations of the quantum group at roots of unity [36, 25, 4]. In section 2, we will give an alternative elementary construction of modular categories producing the same invariants¹.

A skein theoretic construction of the $SU(N)$ invariants was obtained by Yokota [41]; subsequent developments towards the associated TQFT's were made by Lickorish [22, 23]. Our main tool here will be this skein method combined with the structure of the Hecke algebra, which in our context is obtained as the Homflypt skein module of a cylinder $\mathbf{D}^2 \times [0, 1]$ with boundary points. This algebra has been intensively studied (see [12] for a list of references). For our purpose, we emphasize the work of Jones [16] and Wenzl [39]; see also [14], [10]. Our normalization and description of idempotents coincide with those of Aiston-Morton [1, 3, 27].

Following the work of Kirby and Melvin for the $SU(2)$ case [19], Kohno and Takata have studied symmetry formulas for the $SU(N)$ quantum invariants, and defined the $PSU(N)$ invariants [20, 21]. This was used by H. Murakami in [29]. Our reductions and refinements formulas generalize these results.

¹Two modular categories producing the same invariant should be equivalent. A complete treatment of this question has yet to be formed.

Invariants of 3-manifolds from Hecke algebras were obtained by Wenzl in [40]; modular and semi-simple categories from unoriented link invariants (BCD case) are considered by Turaev and Wenzl in [37].

Masbaum and Wenzl [25] have proved the integrality of $SU(N)$ quantum invariants at roots of unity of prime order, and they have shown that this follows from existence of integral modular categories. This is developed by Bruguières in [9].

Acknowledgments. The author wishes to thanks A. Beliakova, A. Bruguières, N. Habegger G. Masbaum, V. Turaev and P. Vogel for useful discussions and suggestions.

1. YOUNG IDEMPOTENTS AND HOMFLYPT SKEIN THEORY

1.1. The Homflypt functor. Let M be an oriented 3-manifold. We denote by $\mathcal{H}(M)$ the k -module freely generated by isotopy classes of framed links in M with the Homflypt relations given in figure 1. Here a framing is a trivialization of the normal bundle; this gives an orientation of the link.

$$\begin{array}{c}
 a^{-1} \begin{array}{c} \nearrow \\ \searrow \end{array} - a \begin{array}{c} \nwarrow \\ \swarrow \end{array} = (s - s^{-1}) \begin{array}{c} \nearrow \\ \searrow \end{array} \begin{array}{c} \nwarrow \\ \swarrow \end{array} \\
 \begin{array}{c} \uparrow \\ \bigcirc \end{array} = av^{-1} \uparrow \qquad \begin{array}{c} \uparrow \\ \bigcirc \end{array} = a^{-1}v \uparrow \\
 L \cup \bigcirc = \frac{v^{-1}-v}{s-s^{-1}} L
 \end{array}$$

FIGURE 1. Homflypt relations

Here k is an integral domain containing the invertible elements a, v, s ; we suppose moreover that $s - s^{-1}$ is invertible in k . (For $L \neq \emptyset$, the third equality is a consequence of the others.)

The Homflypt polynomial [15, 31] gives an isomorphism

$$\begin{array}{ccc}
 \langle \dots \rangle : \mathcal{H}(\mathbf{S}^3) & \rightarrow & k \\
 L & \mapsto & \langle L \rangle
 \end{array}$$

normalized by $\langle \emptyset \rangle = 1$.

An embedding $j : M \hookrightarrow N$ gives a well defined operator $\mathcal{H}(j) : \mathcal{H}(M) \rightarrow \mathcal{H}(N)$. This makes $\mathcal{H}$ into a functor from the category of 3-manifolds whose morphisms are isotopy classes of embeddings, to the category of k -modules.

To an oriented embedding of a disjoint union of solid tori

$$g = \prod_{i=1}^m g_i : \prod_{i=1}^m \mathbf{D}_i^2 \times \mathbf{S}_i^1 \rightarrow \mathbf{S}^3$$

is associated a multilinear map

$$\mathcal{H}(g) : \mathcal{H}(\mathbf{D}^2 \times \mathbf{S}^1)^{\otimes m} \rightarrow \mathcal{H}(\mathbf{S}^3) \approx k$$

This map only depends on the isotopy class of the framed link $L = (L_1, \dots, L_m)$ underlying g . The image of $x_1 \otimes \dots \otimes x_m$ under this map is said to be obtained by cabling the components L_i with the skein elements x_i , and is denoted by $\langle L_1(x_1), \dots, L_m(x_m) \rangle$ or $\langle L(x_1, \dots, x_m) \rangle$.

In the case where M has non-empty boundary, we may fix a framed link l in the boundary of M , and define the relative skein module $\mathcal{H}(M, l)$. By comparing with the orientation of ∂M (our convention for orientation of the boundary is outgoing normal vector first), the framing of l gives an orientation of its components.

1.2. The Hecke category. The k -linear *Hecke category* $\mathbf{H}$ is defined as follows. An object in this category is a disc $\mathbf{D}^2$ equipped with a framed link (note that isotopy is not allowed). If $\alpha = (\mathbf{D}^2, l_0)$ and $\beta = (\mathbf{D}^2, l_1)$ are two objects, the module $\text{Hom}_{\mathbf{H}}(\alpha, \beta)$ is $\mathcal{H}(\mathbf{D}^2 \times [0; 1], l_0 \times 0 \amalg l_1 \times 1)$. The notation $\mathbf{H}(\alpha, \beta)$ and $\mathbf{H}_\alpha$ will be used respectively for $\text{Hom}_{\mathbf{H}}(\alpha, \beta)$ and $\text{End}_{\mathbf{H}}(\alpha)$. For composition, we use the *covariant* notation

$$\begin{aligned} \mathbf{H}(\alpha, \beta) \times \mathbf{H}(\beta, \gamma) &\longrightarrow \mathbf{H}(\alpha, \gamma) \\ (f, g) &\longmapsto fg \end{aligned}$$

When we draw a figure to describe a morphism, the *time* parameter goes upwards, so that the morphism fg is depicted with g lying above f .

Note. The terminology *Hecke category* was introduced by Turaev in [34]. His definition gives a category equivalent to an unframed version of ours.

We will use the notation $\hat{f}$ for the closure in $\mathcal{H}(\mathbf{D}^2 \times \mathbf{S}^1)$ of the morphism $f \in \mathbf{H}_\alpha$. Let U_0 be a 0-framed unknot in $\mathbf{S}^3$. We use the notation $\langle f \rangle$ for $\langle U_0(\hat{f}) \rangle$ (the quantum trace).

We will simply denote by n the object formed by n standard points (along the real axis) equipped with the standard framing $(1, \sqrt{-1})$.

By using a standard embedding $\mathbf{D}^2 \amalg \mathbf{D}^2 \hookrightarrow \mathbf{D}^2$, we make $\mathbf{H}$ into a monoidal category². This category has a braiding and a twist operator.

²The associativity isomorphisms just move the embedded discs along the real axis, and will be omitted. Note that the object $1^{\otimes n}$ is defined up to these associativity isomorphisms, and is canonically isomorphic to the standard object n .

We can define in $\mathcal{H}$ a duality rule so that we get a ribbon category ([35], see also [38]). We proceed as follows. To an object $\alpha = (\mathbf{D}^2, l)$, we associate the object $\alpha^* = (\mathbf{D}^2, -\bar{l})$, where $-\bar{l}$ is the framed link obtained by applying to l the differential of $z \mapsto -\bar{z}$, and define the morphisms $b_\alpha \in H(0, \alpha \otimes \alpha^*)$ and $d_\alpha \in H(\alpha^* \otimes \alpha, 0)$ according to the figure below (a copy of 1_α is embedded along the framed arc).

$$b_\alpha = \alpha \curvearrowright \quad d_\alpha = \curvearrowleft \alpha$$

Our purpose is to discuss which modular categories arise from this ribbon Hecke category. For this, we need a finite set of simple objects with nice properties ([35] p74).

1.3. Idempotents in the Hecke algebra. The algebra H_n is isomorphic to the quotient of the algebra of the braid group $k[B_n]$ by the Homflypt relation

$$a^{-1} \begin{array}{c} \nearrow \\ \searrow \end{array} - a \begin{array}{c} \nwarrow \\ \nearrow \end{array} = (s - s^{-1}) \begin{array}{c} \nearrow \\ \nearrow \end{array} \begin{array}{c} \searrow \\ \searrow \end{array}$$

which is the Hecke algebra of type A_{n-1} . An unframed version of this result was proved independently by Morton-Traczyk [28] and Turaev [34]. This algebra is known to be generically semi-simple. It is a deformation of the algebra of the symmetric group, and its structure can be obtained by extending the classical Young theory [8, 16, 39, 14, 10].

Recall that, to a partition of n , $\lambda = (\lambda_1 \geq \dots \geq \lambda_p \geq 1)$, $\lambda_1 + \dots + \lambda_p = n$, is associated a Young diagram of size $|\lambda| = n$, which we denote also by λ . This diagram has n cells indexed by $\{(i, j), 1 \leq i \leq p, 1 \leq j \leq \lambda_i\}$.

If c is the cell of index (i, j) in a Young diagram, its hook-length $hl(c)$ and its content $cn(c)$ are defined by

$$hl(c) = \lambda_i + \lambda_j^\vee - i - j + 1, \quad cn(c) = j - i.$$

Here $\lambda^\vee$ is the transposed Young diagram and $\lambda_j^\vee$ is the length of the j -th column of λ (the j -th line of $\lambda^\vee$).

For $n \geq 0$, the quantum integer $[n]$, and the quantum factorial $[n]!$ are defined by $[n] = \frac{s^n - s^{-n}}{s - s^{-1}}$ and $[n]! = \prod_{j=1}^n [j]$.

For a Young diagram λ , we will use the notation $[hl(\lambda)]$, for the product over all cells of the quantum hook-lengths.

$$[hl(\lambda)] = \prod_{\text{cells}} [hl(c)]$$

Symmetrizers. Let $\sigma_i \in H_n$, $i = 1, \dots, n-1$, be represented by the standard generators of the braid group B_n (the strand numbered i crosses over the strand numbered $i+1$).

Proposition 1.1. *If $[n]!$ is invertible in k , then there exists a unique idempotent $f_n \in H_n$ such that $\forall i \sigma_i f_n = a s f_n = f_n \sigma_i$, and a unique idempotent $g_n \in H_n$ such that $\forall i \sigma_i g_n = -a s^{-1} g_n = g_n \sigma_i$.*

Proof. It is shown in [26, 3] that the deformation f_n of the Young symmetrizer, given below, satisfies the required condition.

$$f_n = \frac{1}{[n]!} s^{-\frac{n(n-1)}{2}} \sum_{\pi \in S_n} (a s^{-1})^{-l(\pi)} w_\pi$$

Here w_π is the positive braid associated with the permutation π , and $l(\pi)$ is the length of π .

One can also construct f_n recursively using the formulas below ([41]).

$$[2]f_2 = s^{-1} \left(\begin{array}{c} \nearrow \\ \searrow \end{array} \right) + a^{-1} \left(\begin{array}{c} \nwarrow \\ \searrow \end{array} \right)$$

$$[n+1]f_{n+1} = -[n-1]f_n \otimes 1_1 + [2][n](f_n \otimes 1_1)(1_{n-1} \otimes f_2)(f_n \otimes 1_1)$$

(Here, $1_p \in H_p$ is the identity.)

Suppose now that f'_n satisfies the condition of the theorem. Then we have that $f_n f'_n = f'_n = \zeta f_n$, and $\zeta = 1$ by idempotence.

For g_n we can proceed similarly, either with the deformed antisymmetrizer

$$g_n = \frac{1}{[n]!} s^{\frac{n(n-1)}{2}} \sum_{\pi \in S_n} (-a s)^{-l(\pi)} w_\pi$$

or with the recursive formulas

$$g_1 = 1_1$$

$$g_{n+1} = 1_1 \otimes g_n - \frac{[2][n]}{[n+1]} (1_1 \otimes g_n)(f_2 \otimes 1_{n-1})(1_1 \otimes g_n)$$

□

Note that $1_2 = f_2 + g_2$; this can be used to obtain more symmetric recursive formulas for f_n and g_n . Our choice immediately gives the following lemma [41], which is useful in Yokota's skein computations.

Lemma 1.1. *For any two integers p, q such that $[p+1]!$ and $[q+1]!$ are invertible, one has*

$$\begin{aligned} [p+q]f_p \otimes g_q &= [p+1][q](1_p \otimes g_q)(f_{p+1} \otimes 1_{q-1})(1_p \otimes g_q) \\ &\quad + [p][q+1](f_p \otimes 1_q)(1_{p-1} \otimes g_{q+1})(f_p \otimes 1_q) \end{aligned}$$

Aiston-Morton description of Young symmetrizers. For a Young diagram λ of size n , we denote by $\square_\lambda$ the object of the category $\mathbf{H}$ formed with one point for each cell c of λ ; if c has index (i, j) (i -th line, and j -th column), then the corresponding point in $\mathbf{D}^2$ is $\frac{j+i\sqrt{-1}}{n}$. Suppose that $\lambda = (\lambda_1 \geq \dots \geq \lambda_p \geq 1)$, and that $\lambda^\vee = (\lambda_1^\vee \geq \dots \geq \lambda_q^\vee \geq 1)$ is the transposed Young diagram.

Let F_λ (resp. G_λ) be the element in $\mathbf{H}_{\square_\lambda}$ formed with one copy of $[\lambda_i]!f_{\lambda_i}$ along the line i , for $i = 1, \dots, p$ (resp. one copy of $[\lambda_j^\vee]!g_{\lambda_j^\vee}$ along the column j , for $j = 1, \dots, q$). Note that these expressions have no denominators.

In the proposition below, $<$ denotes the lexicographic ordering, and λ and μ are Young diagrams with the same number of cells.

Proposition 1.2. *a) If $\mu < \lambda$, then $F_\lambda \mathbf{H}(\square_\lambda, \square_\mu) G_\mu = 0$.
b) If $\lambda < \mu$, then $G_\lambda \mathbf{H}(\square_\lambda, \square_\mu) F_\mu = 0$.
c) One has $F_\lambda \mathbf{H}_{\square_\lambda} G_\lambda = k F_\lambda G_\lambda$.*

This proposition is proved in [3], section 4. We outline here the proof given there.

Proof. a) Every family of $|\lambda|!$ braids in $\mathbf{H}(\square_\lambda, \square_\mu)$, which induces all bijections between the cells of λ and the cells of μ , is a basis of $\mathbf{H}_{\square_\lambda}$. The hypothesis $\mu < \lambda$ implies that any bijection between the cells of λ and the cells of μ carries at least two cells in some line of λ , to cells in the same column of μ . So we can find a basis of $\mathbf{H}(\square_\lambda, \square_\mu)$ represented by braids which connect in a separate cylinder two points in some line of $\square_\lambda$ with two points in some column of $\square_\mu$. One can deduce that $F_\lambda \mathbf{H}(\square_\lambda, \square_\mu) G_\mu = 0$. Assertion b) is shown similarly.

c) We say that a permutation π between the cells of λ does not separate, if some pair of cells in the same line is mapped to some pair of cells in the same column; other permutations are said to separate. Using that any separating permutation π of the cells can be written $\pi = \pi_R \pi_C$ (permutations act on the left), where π_R (resp. π_C) preserves the rows (resp. the columns), we see that we can find a basis represented by braids b_π indexed by permutations such that

$$\begin{cases} F_\lambda b_\pi G_\mu = 0 & \text{if } \pi \text{ does not separate,} \\ F_\lambda b_\pi G_\mu = \zeta_\pi F_\lambda G_\mu & \text{if } \pi \text{ separates.} \end{cases}$$

Statement c) follows. $\square$

The argument used in the proof of a) above shows the following.

Lemma 1.2. *a) If $p > \lambda_1$, and for some object α , $x = x_1 \otimes f_p \otimes x_2 \in \mathbf{H}_\alpha$, then $x \mathbf{H}(\alpha, \square_\lambda) G_\lambda = 0$.*

b) If $p > \lambda_1^\vee$, and for some object α , $x = x_1 \otimes g_p \otimes x_2 \in H_\alpha$, then $F_\lambda H(\square_\lambda, \alpha)x = 0$.

Let $\tilde{y}_\lambda = F_\lambda G_\lambda$. A consequence of proposition 1.2 is that $\tilde{y}_\lambda$ is a quasi-idempotent.

Proposition 1.3. *One has $\tilde{y}_\lambda^2 = [hl(\lambda)]\tilde{y}_\lambda$.*

This proposition is proved by Yokota in [41], lemma 2.3 (see also [2]); the fact that our parameters a and v are more generic is irrelevant in this skein computation.

If $[hl(\lambda)]$ is invertible, we define the idempotent y_λ by

$$y_\lambda = [hl(\lambda)]^{-1}\tilde{y}_\lambda$$

Suppose that λ and μ are Young diagrams with the same number of cells, such that $[hl(\lambda)]$ and $[hl(\mu)]$ are invertible. From proposition 1.2, we have.

Proposition 1.4. a) If $\mu \neq \lambda$, then $y_\lambda H(\square_\lambda, \square_\mu)y_\mu = 0$.
b) One has that $y_\lambda H_{\square_\lambda} y_\lambda = ky_\lambda$.

Let $\lambda \subset \mu$ be two Young diagrams, the complement of λ in μ is called a skew Young diagram and is denoted by μ/λ . As above we can define the object $\square_{\mu/\lambda}$ in the category H with one point for each cell. The following is proven in the same way as proposition 1.3.

Lemma 1.3. *Let $\lambda \subset \mu$ be two Young diagrams, and let $v \in H_{\square_{\mu/\lambda}}$. One has that*


$$F_\mu(\tilde{y}_\lambda \otimes v)G_\mu = [hl(\lambda)]F_\mu(1_{\square_\lambda} \otimes v)G_\mu$$

We will need the following formulas.

Proposition 1.5. a) Suppose that $\lambda \subset \mu$ are two Young diagrams such that the skew diagram μ/λ has only one cell c ($|\mu| = |\lambda| + 1$). Then

$$= a^{2|\lambda|} s^{2cn(c)} y_\mu$$

b) (Framing coefficient)



$$= a^{|\lambda|^2} v^{-|\lambda|} s^{2 \sum_{cells} cn(c)} y_\lambda$$

Statements a) and b) can be deduced from each other. Statement b) is theorem 5.5 of [3]; the proof given there is an elementary skein calculation which contains statement a). These results can also be deduced from [40]. Statement b) is a framed version of lemma 3.2.1 there.

1.4. Structure of the generic Hecke algebra and path idempotents. Here we suppose that n is fixed, and that, in the domain k , $[j]$ is invertible for every $j \leq n$, so that all the idempotents y_λ exist in H_n .

A standard tableau t with shape a Young diagram $\lambda = \lambda(t)$ is a labelling of the cells, with the integers 1 to n , which is increasing along lines and columns. We denote by t' the tableau obtained by removing the cell numbered by n . We define $\alpha_t \in H(n, \square_\lambda)$ and $\beta_t \in H(\square_\lambda, n)$ by

$$\begin{aligned} \alpha_1 &= \beta_1 = 1_1 \\ \alpha_t &= (\alpha_{t'} \otimes 1_1) \varrho_t y_\lambda \\ \beta_t &= y_\lambda \varrho_t^{-1} (\beta_{t'} \otimes 1_1) \end{aligned}$$

Here $\varrho_t \in H(\square_{\lambda(t')} \otimes 1, \square_\lambda)$ is a standard isomorphism which is obtained by moving the added point, first along a line parallel to the imaginary axis, and then along a line parallel to the real axis.

Note that $\beta_\tau \alpha_t = 0$ if $\tau \neq t$, and $\beta_t \alpha_t = y_{\lambda(t)}$.

Proposition 1.6. *The family $\alpha_t \beta_\tau$ for all standard tableaux t, τ such that $\lambda(t) = \lambda(\tau)$ forms a basis for H_n .*

Proof. We have (here δ is the Kronecker delta).

$$\alpha_t \beta_\tau \alpha_s \beta_\sigma = \delta_{\tau s} \alpha_t \beta_\sigma$$

This shows independence. Moreover the number of vectors is equal to the dimension. This shows the proposition over the field of quotients of the domain k . The above formula gives the coordinate forms and shows that the result is valid over k . $\square$

This gives explicitly the semi-simple decomposition of H_n . The simple components are indexed by Young diagrams. The component indexed by λ is the two-sided ideal generated by y_λ ; its rank is d_λ^2 , where d_λ is the number of standard tableaux with shape λ . The $\alpha_t \beta_\tau$ are *matrix units*

in the sense of Ram-Wenzl [32]. The diagonal elements $p_t = \alpha_t \beta_t$ are the path idempotents described in [39]; the minimal central idempotents are the $z_\lambda = \sum_{\lambda(t)=\lambda} p_t$.

Theorem 1.1 (Branching formula).

$$y_\lambda \otimes 1_1 = \sum_{\substack{\lambda \subset \mu \\ |\mu| = |\lambda| + 1}} (y_\lambda \otimes 1_1) y_\mu (y_\lambda \otimes 1_1)$$

We have omitted in this formula the standard isomorphism between $\lambda \otimes 1$ and μ .

This branching formula for the path idempotents is given by Wenzl in [39]. A similar formula is given by Yokota [41], prop. 2.11, with a proof using skein calculus.

Proof. We continue to denote by z_μ the minimal central idempotent corresponding to the simple component indexed by μ in $H_{\square_\lambda \otimes 1}$. We have

$$y_\lambda \otimes 1_1 = \sum_{\mu} z_\mu (y_\lambda \otimes 1_1) = \sum_{\substack{\lambda \subset \mu \\ |\mu| = |\lambda| + 1}} (y_\lambda \otimes 1_1) z_\mu (y_\lambda \otimes 1_1)$$

From the formula $z_\mu = \sum_{\lambda(t)=\mu} p_t$, we see that in the above, only those tableaux t with $\lambda(t') = \lambda$ contribute, and all these contributions are proportional to $(y_\lambda \otimes 1_1) y_\mu (y_\lambda \otimes 1_1)$, by lemma 1.3. By using idempotence, we have that $(y_\lambda \otimes 1_1) z_\mu (y_\lambda \otimes 1_1) = (y_\lambda \otimes 1_1) y_\mu (y_\lambda \otimes 1_1)$. $\square$

Proposition 1.7 (Quantum dimension).

$$\langle y_\lambda \rangle = \prod_{\text{cells}} \frac{v^{-1} s^{cn(c)} - v s^{-cn(c)}}{s^{hl(c)} - s^{-hl(c)}}$$

Recall that $\langle y_\lambda \rangle$ is the Homflypt polynomial of a 0-framed unknot cabled with the closure $\hat{y}_\lambda$ of y_λ in $\mathbf{D}^2 \times \mathbf{S}^1$.

We will denote $\langle y_\lambda \rangle$ simply by $\langle \lambda \rangle$, and call it the quantum dimension of λ .

The assertion above can be proven by a skein calculation (see proposition 2.4 in [41] or [2]). An alternative proof ([39]) is to use the Young algebra. From the branching formula (1.1), we can deduce that $\hat{y}_\lambda$ is the λ -indexed Schur polynomial in the $\hat{g}_k$; hence the general formula follows from $\langle g_k \rangle = \prod_{j=1}^k \frac{v^{-1} s^{j-1} - v s^{1-j}}{s^j - s^{-j}}$, which can be proven by using the recursive formula for g_k .

1.5. The $\mathcal{C}$ -completed Hecke category. Suppose $\mathcal{C}$ is a set of Young diagrams λ , such that the Young idempotents y_λ exist. The $\mathcal{C}$ -completed Hecke category $H^\mathcal{C}$ is defined as follows.

An object in this category is a disc $\mathbf{D}^2$ equipped with a labelled framed link $l = ((l_1, \lambda^{(1)}), \dots, (l_m, \lambda^{(m)}))$, where $\lambda^{(1)}, \dots, \lambda^{(m)}$ are Young diagrams in $\mathcal{C}$.

If $\alpha = (\mathbf{D}^2, l)$ is such an object, its expansion $E(\alpha) = (\mathbf{D}^2, E(l))$ is obtained by embedding the object $\square_\lambda$ in a neighbourhood of l_i , according to the framing. The tensor product $y_{\lambda^{(1)}} \otimes \dots \otimes y_{\lambda^{(m)}}$ defines an idempotent $\pi_\alpha \in H_{E(\alpha)}$. The module $H^\mathcal{C}(\alpha, \beta)$ is defined by

$$H^\mathcal{C}(\alpha, \beta) = \pi_\alpha H(E(\alpha), E(\beta)) \pi_\beta$$

The duality extends to the category $H^\mathcal{C}$ in a natural way, and we again have a ribbon category. We denote simply by λ the object of $H^\mathcal{C}$ which is a disc $\mathbf{D}^2$ with the origin labelled by λ .

1.6. Homflypt calculus using ribbon graphs. Following Turaev ([35], I.2), we can define the category Rib_H of ribbon graphs over H , and use the canonical functor $F_H : Rib_H \rightarrow H$. A colored ribbon graph gives a morphism in the category H (the functor F_H will be implicit). This can be described as follows. For each band in the graph, colored with α , embed (using the framing of the band) a copy of 1_α , the identity of α ; for each loop colored with α embed a copy of $\hat{1}_\alpha$, the closure of 1_α ; for each coupon colored with the morphism f , embed a copy of f .

We can proceed similarly with the $\mathcal{C}$ -completed Hecke category $H^\mathcal{C}$.

2. THE MODULAR CATEGORIES $H^{SU(N,K)}$ AND $H^{PSU(N,K)}$

2.1. Roots of unity. In this section, we suppose that s is a primitive $2(N+K)$ -th root of unity, and that $v = s^{-N}$ (rank N and level K , $N \geq 2$ and $K \geq 1$). We suppose moreover that $N+K$ is invertible in k . A consequence is that $[n]$ is invertible for $n < N+K$. This can be seen as follows. We have that

$$[n] = s^{-n+1} \prod_{1 \leq j \leq n} \phi_j(s^2)$$

Here $\phi_j \in \mathbb{Z}[X]$ is the j -indexed cyclotomic polynomial. The required invertibility is a consequence of the following lemma. Here we suppose $p \geq 2$.

Lemma 2.1. *If $j \notin p\mathbb{Z}$, then ϕ_j divides p in $\mathbb{Z}[X]/\phi_p$.*

Proof. Let $d = \gcd(p, j)$. In $\mathbb{Z}[X]$, one has the relation $U(X^j - 1) + V(X^p - 1) = X^d - 1$. If $j > d$, this implies the relation $U_1 \phi_j + V_1 \phi_p = 1$,

hence ϕ_j is invertible in $\mathbb{Z}[X]/\phi_p$. If $j = d$ then using the derivative of $X^p - 1 = \phi_j T$, we get $p \equiv XT'\phi_j \bmod \phi_p$. $\square$

We observe that the idempotent y_λ exists for every Young diagram λ with $\lambda_1 + \lambda_1^\vee \leq N + K$. We denote by $\mathcal{C}_{N,K}$ the set of these Young diagrams and consider the category $H^{(N,K,a)}$ obtained from the $\mathcal{C}_{N,K}$ -completed Hecke category, $H^{\mathcal{C}_{N,K}}$ by applying the following purification procedure ([35] p504).

Definition 2.1. A morphism $f \in H^{\mathcal{C}_{N,K}}(\alpha, \beta)$ is negligible iff

$$\forall g \in H^{\mathcal{C}_{N,K}}(\beta, \alpha), \quad \langle fg \rangle = 0$$

Objects of $H^{(N,K,a)}$ are those of $H^{\mathcal{C}_{N,K}}$. The module $H^{(N,K,a)}(\alpha, \beta)$ is the quotient of $H^{\mathcal{C}_{N,K}}(\alpha, \beta)$ by the submodule of negligible morphisms. One can verify that composition and tensor product are well defined on the quotient. The negligible morphisms give local relations in Homflypt modules. We denote by $\mathcal{H}^{(N,K,a)}(M)$, the quotient of $\mathcal{H}^{\mathcal{C}_{N,K}}(M)$ by these relations. Note that for $M = \mathbf{S}^3$, no new relation appears. In the case $M = \mathbf{D}^2 \times \mathbf{S}^1$, the algebra structure is well defined on the quotient. The multilinear form $\langle L(\dots) \rangle$ is well defined on $\mathcal{H}^{(N,K,a)}(\mathbf{D}^2 \times \mathbf{S}^1)$. More generally, ribbon graphs may be colored using the category $H^{(N,K,a)}$.

For a Young diagram $\lambda = (\lambda_1, \dots, \lambda_p) \in \mathcal{C}_{N,K}$, the identity morphism 1_λ is negligible if and only if its quantum dimension $\langle \lambda \rangle$ is zero. This will be the case if and only if $K < \lambda_1$ or $N < \lambda_1^\vee$. We will use the following sets of Young diagrams.

$$\bar{\Gamma}_{N,K} = \{(\lambda_1, \dots, \lambda_p), \lambda_1 \leq K \text{ and } p \leq N\}$$

$$\Gamma_{N,K} = \{(\lambda_1, \dots, \lambda_p), \lambda_1 \leq K \text{ and } p < N\}$$

Denote by 1^N (resp. (K)) the diagram, with one column containing N cells (resp. one line containing K cells). The proposition below shows that 1^N (resp. (K)) is practically influential. It will also explain our choice of the framing parameter a .

Proposition 2.1. *The following identities hold in the category $H^{(N,K,a)}$*

$$\begin{array}{c} \uparrow \\ \text{---} \boxed{g_N} \text{---} \uparrow \end{array} = a^{2N} s^2 \begin{array}{c} \uparrow \\ \text{---} \boxed{g_N} \text{---} \uparrow \end{array}$$

$$\begin{array}{c} \uparrow \\ \text{---} \downarrow \uparrow \end{array} \boxed{f_K} = a^{2K} s^{-2} \begin{array}{c} \uparrow \uparrow \\ \text{---} \end{array} \boxed{f_K}$$

Proof. We justify the first equality. The proof of the second one is similar. The branching formula 1.1 gives (g_{N+1} is negligible)

$$g_N \otimes 1_1 = (g_N \otimes 1_1) y_{(2, 1^{N-1})} (g_N \otimes 1_1)$$

The formula comes from proposition 1.5. $\square$

Observe that (at rank N , level K), we have $\langle g_N \rangle = 1$, and that the framing coefficient for g_N is $(a^N s)^N$. Using this we obtain the following corollary.

Corollary 2.1. *Suppose that $a^N s = 1$. Then the morphism represented by a colored ribbon graph is not changed*

*if some band colored with the object 1^N is twisted or moved
across any other band,
or if some loop colored with the object 1^N is removed.*

Until the end of this section, we suppose that $a^N s = 1$. If λ is a Young diagram in $\bar{\Gamma}_{N,K}$, we denote by λ^* the skew diagram μ/λ , with $\mu = \lambda_1^N$; up to a rotation in the plane, λ^* is a Young diagram in $\Gamma_{N,K}$.

Lemma 2.2. *a) For any $\lambda \in \bar{\Gamma}_{N,K}$, one has $\langle \lambda^* \rangle = \langle \lambda \rangle$.*

b) For any $\lambda \in \bar{\Gamma}_{N,K}$ of the form $\lambda = 1^N + \nu$, one has $\langle L(\hat{y}_\lambda, \dots) \rangle = \langle L(\hat{y}_\nu, \dots) \rangle$.


In particular, for $j \leq K$, one has $\langle j^N \rangle = 1$

Proof. If $v = s^{-N}$ (rank N), the quantum dimension is given by

$$\langle \lambda \rangle = \prod_{\text{cells}} \frac{[N + cn(c)]}{[hl(c)]}.$$

$\square$

Lemma 2.3. *For any $\lambda \in \bar{\Gamma}_{N,K}$, we have*



$$= \frac{1}{\langle \lambda \rangle} y_\lambda.$$

Proof. From the definition of the idempotents, we see that the left hand side is proportional to y_λ . The coefficient is obtained by comparing the quantum traces. $\square$

Reversing orientation. Let L be a framed link, and let L' be the framed link obtained from L by reversing the orientation of the first component (i.e. by changing the sign of the second vector in the trivialization of the normal bundle).

Proposition 2.2. *For any $\lambda \in \bar{\Gamma}_{N,K}$, one has*

$$\langle L(\hat{y}_\lambda, \dots) \rangle = \langle L'(\hat{y}_{\lambda^*}, \dots) \rangle$$

Proof. Using lemma 2.3, we introduce somewhere the idempotent associated with the diagram λ_1^N . Then we can cut the band colored with λ_1^N , move one of the ends along the first component L_1 of L and glue it back again, so that the band colored with λ^* goes along L_1 with the reverse orientation. By corollary 2.1 this does not change the evaluated Homflypt polynomial. Using lemma 2.3 again, we get the result. $\square$

The above suggests that we could build an isomorphism between the trivial object 0 and 1^N , and also between the dual of λ (denoted by λ^*) and λ^* (note the difference between $*$ and $\star$). In order to do that, we will add some morphisms.

We saw that we can define morphisms in the Hecke category, and more generally skein elements in Homflypt skein modules by using colored ribbon graphs. We extend the $\mathcal{C}_{N,K}$ -completed Hecke category and the Homflypt skein theory by allowing incoming or outgoing vertices, colored with $r = 1^N$. Together with the relation of isotopy rel. boundary, the Homflypt relations, and the negligible morphisms, we add the relation given by gluing an incoming r -colored vertex with an outgoing one. (Note that the half twist is not trivial; hence the orientations of the glued ends of the ribbon edges must be respected.)

We will denote by $\overline{H}^{(N,K,a)}$ the category whose objects are those of $H^{(N,K,a)}$ and whose morphisms are defined using the extended Homflypt skein theory above, and by $\overline{\mathcal{H}}^{(N,K,a)}$ the extended Homflypt skein functor.

From corollary 2.1, we deduce that the Homflypt polynomial extends to an isomorphism $\overline{\mathcal{H}}^{(N,K,a)}(\mathbf{S}^3) \approx k$.

In the category $\overline{H}^{(N,K,a)}$, an object $\lambda \in \Gamma_{N,K}$ is still simple, and its dual λ^* is isomorphic to λ^* .

From the branching formula (theorem 1.1), we can see that the simple objects in $\Gamma_{N,K}$ dominate the category $\overline{H}^{(N,K,a)}$. This gives us all the defining properties of a modular category except the non-degeneracy axiom. We say that $\overline{H}^{(N,K,a)}$ is a *pre-modular category*.

Note that distinct objects $\lambda, \mu \in \Gamma_{N,K}$ are not isomorphic, but this does not imply the non-degeneracy axiom.

2.2. The handle slide condition. We say that $\Omega \in \mathcal{H}^{(N,K,a)}(\mathbf{D}^2 \times \mathbf{S}^1)$ satisfies the Kirby condition if

$$(K) \quad \begin{cases} \forall x \in \mathcal{H}^{(N,K,a)}(\mathbf{D}^2 \times \mathbf{S}^1) \quad \langle H_1(x, \Omega) \rangle = \langle U_0(x) \rangle \langle U_1(\Omega) \rangle \\ \langle U_1(\Omega) \rangle \text{ is invertible} \end{cases}$$

Here, for $\epsilon \in \{-1, 0, 1\}$ we denote by U_ϵ the unknot with framing ϵ , and by H_ϵ the Hopf link with linking number one and both components having framing ϵ .

A solution of the above is essentially unique. In [6], we discussed this condition in the context of a formal skein theory.

A framed link L determines by surgery a 3-manifold denoted by $S^3(L)$ (every compact oriented 3-manifold can be obtained in this way). As a consequence of Kirby's theorem [18], if a solution Ω exists, then an appropriate normalization of $\langle L(\Omega, \dots, \Omega) \rangle$ is an invariant of the surgered manifold $M = \mathbf{S}^3(L)$. It is convenient to choose a solution ω of (K) such that $\langle U_1(\omega) \rangle \langle U_{-1}(\omega) \rangle = 1$. (It may be necessary to extend k by adding a square root.) An appropriate normalization is then

$$\tau(M) = \langle U_1(\omega) \rangle^{-\sigma(L)} \langle L(\omega, \dots, \omega) \rangle$$

Here $\sigma(L)$ is the signature of the linking matrix, i.e. $\sigma(L) = b_+ - b_-$, where b_+ (resp. b_-) is the number of positive (resp. negative) eigenvalues of the linking matrix B_L associated with L .

Recall that we have denoted by $\Gamma_{N,K}$ the set of Young diagrams with at most $N - 1$ lines and K columns (the empty diagram is included).




FIGURE 2. Sliding property

We set

$$\Omega = \sum_{\lambda \in \Gamma_{N,K}} \langle \lambda \rangle \hat{y}_\lambda$$

Following Yokota [41], we can show the sliding property. From this sliding property and the proposition 2.2 we can deduce the first part of the Kirby condition (the handle slide).

Proposition 2.3 (Sliding property). *The Homflypt polynomial of a link in $\mathbf{S}^3$, which has one of its components cabled with the skein element Ω , satisfies the equality in figure 2.*

In this figure, the dashed curve means that the component cabled with Ω may be non-trivially embedded in the sphere; the bracket notation for the Homflypt polynomial is omitted.

Proof. Using the definition of Ω , the ramification formula, and the lemma 2.3.

$$lhs = \sum_{\lambda} \langle \lambda \rangle \sum_{\substack{\lambda \subset \mu \\ |\mu| = |\lambda| + 1}} \langle \mu \rangle \langle A(\lambda, \mu) \rangle,$$

where $A(\lambda, \mu)$ is the skein element represented in figure 3. Note that in the above, we can forget any μ with $\mu_1 = K + 1$, because in this case the corresponding skein element contains a negligible morphism. We can cut the band colored with μ_1^N , move one of the ends along the closed component and glue it back again, so that the band colored with μ^* goes along this closed component with the reverse orientation. Then we apply lemma 2.3 and the ramification formula again. Using the involution $\star$ on the set $\Gamma_{N,K}$, and also lemma 2.2, we obtain the required equality. $\square$

2.3. The invariant $\tau^{SU(N,K)}$, and the modular category $H^{SU(N,K)}$. Until the end of section 2, we will consider the choice of the framing parameter $a = s^{\frac{-1}{N}}$. More precisely a is a primitive $2N(K + N)$ -th root




FIGURE 3. The rectangle trick

of unity, and $s = a^{-N}$.

For this choice of parameter a , we denote the category $\overline{\mathbf{H}}^{(N,K,a)}$ by $\mathbf{H}^{SU(N,K)}$, and the corresponding skein functor by $\mathcal{H}^{SU(N,K)}$.

Lemma 2.4.

$$\langle U_1(\Omega) \rangle \langle U_{-1}(\Omega) \rangle = \langle \Omega \rangle = (-1)^{\frac{N(N-1)}{2}} \frac{N(N+K)^{N-1}}{\prod_{1 \leq j \leq N} (s^j - s^{-j})^{2(N-j)}}$$

Proof. The first equality comes from the sliding property and lemma 2.5 below. For the second equality, we follow Erlijman's computation in [11].

$$\langle \lambda \rangle = \frac{\prod [N + cn]}{\prod [hl]} = s^{-(N-1)|\lambda|} \mathcal{S}_\lambda(1, s^2, \dots, s^{2(N-1)})$$

Here $\mathcal{S}_\lambda$ is the λ -indexed Schur symmetric polynomial (see ch1 in [24]),

$$\langle \lambda \rangle^2 = \frac{a_{\rho+\lambda} \bar{a}_{\rho+\lambda}}{a_\rho \bar{a}_\rho}$$

with $\rho = (N-1, N-2, \dots, 0)$ and, for $l = (l_1, \dots, l_N)$

$$a_l = \det(s^{2(i-1)l_j})_{1 \leq i, j \leq N}$$

We have

$$\begin{aligned} a_\rho \bar{a}_\rho &= \prod_{i < j} (s^{2j} - s^{2i})(s^{-2j} - s^{-2i}) = (-1)^{\frac{N(N-1)}{2}} \prod_{\nu=1}^{N-1} (s^\nu - s^{-\nu})^{2(N-\nu)} \\ \sum_{\lambda \in \Gamma_{N,K}} a_{\rho+\lambda} \bar{a}_{\rho+\lambda} &= \sum_{N+K > l_1 > \dots > l_N = 0} \det(s^{2(i-1)l_j}) \det(s^{-2(i-1)l_j}) \\ \sum_{\lambda \in \Gamma_{N,K}} a_{\rho+\lambda} \bar{a}_{\rho+\lambda} &= \frac{1}{(N-1)!} \sum_{0 \leq l_1, \dots, l_{N-1} < N+K} \sum_{\pi, \pi' \in \mathcal{S}_N} \epsilon_\pi \epsilon_{\pi'} \prod_i s^{2l_i(\pi(i) - \pi'(i))} \end{aligned}$$

Here, we have symmetrized the index set for l (and added terms which are zero), and we expanded the determinants.


$$\sum_{\lambda \in \Gamma_{N,K}} a_{\rho+\lambda} \bar{a}_{\rho+\lambda} = \frac{1}{(N-1)!} \sum_{\pi, \pi'} \epsilon_{\pi} \epsilon_{\pi'} \prod_{i=1}^{N-1} \sum_{l_i=0}^{N+K-1} s^{2l_i(\pi(i)-\pi'(i))}$$

For $\pi \neq \pi'$, a zero term appears, and the $N!$ remaining terms are all equal. We get


$$\sum_{\lambda \in \Gamma_{N,K}} a_{\rho+\lambda} \bar{a}_{\rho+\lambda} = N(N+K)^{N-1}$$

Our formula follows. $\square$

Lemma 2.5. *If $\lambda \neq \emptyset$, then the following morphism is zero in $H^{SU(N,K)}$.*



Proof. By the sliding property, for any Young diagram μ such that $\lambda \subset \mu$, and $|\mu/\lambda| = 1$, we have



By proposition 1.5, if the color λ is not killed by inserting a 0-framed meridian cabled with Ω , then $a^{2|\lambda|} s^{2cn(\mu/\lambda)} = 1$ for any μ as above with $\langle \mu \rangle \neq 0$. If two such μ exist, with added cells of respective indices (i, j) and (i', j') , then $s^{2(j'-j+i-i')} = 1$ and $-(N+K) < j'-j+i-i' < N+K$. This implies $(i, j) = (i', j')$. The remaining diagrams are those with K columns and n lines, $n < N$. In this case we get $a^{2nK} s^{-2n} = a^{2n(K+N)} = 1$. The order of a implies $n = 0$. $\square$

We suppose that $N(N+K)$ is invertible. We set $\eta^{-2} = \langle \Omega \rangle$ (we extend k if necessary), $\omega = \eta\Omega$, $\Delta = \langle U_1(\omega) \rangle$.

Theorem 2.1. *There exists an invariant of compact oriented 3-manifolds defined on a surgery presentation by the following formula.*

$$\tau^{SU(N,K)}(\mathbf{S}^3(L)) = \Delta^{-\sigma(L)} \langle L(\omega, \dots, \omega) \rangle$$

This invariant can be extended to manifolds with (colored) links by the formula

$$\tau^{SU(N,K)}(\mathbf{S}^3(L), K) = \Delta^{-\sigma(L)} \langle L(\omega, \dots, \omega) \cup K \rangle$$

The above is a consequence of the Kirby theorem. Using [35], the following constructs the associated TQFT (and proves again the theorem above).

Theorem 2.2. *The category $\mathcal{H}^{SU(N,K)}$ is a modular category with $\Gamma_{N,K}$ as a representative set of isomorphism classes of simple objects.*

Proof. We have to check invertibility of the S matrix whose entries, indexed by $\Gamma_{N,K}$, are the evaluations of the Homflypt polynomial for a Hopf link H_0 (0-framed, and with linking +1), whose components are colored with the corresponding indices. The lemma below gives the result (the matrix $\overline{S}$ is the conjugate of S). $\square$

Lemma 2.6. *One has (I is the unit matrix)*

$$S\overline{S} = \langle \Omega \rangle I$$

Proof. The (λ, μ) -indexed entry of the matrix $S\overline{S}$ can be written

$$u_{\lambda\mu} = \langle H_0(\hat{y}_\lambda \hat{y}_{\mu^*}), \Omega \rangle$$

We write $u_{\lambda\mu}$ as the quantum trace of the morphism $\gamma_{\lambda\mu}$ represented by a $\lambda \otimes \mu^*$ -colored band together with an Ω -cabled meridian. Using that the objects in $\Gamma_{N,K}$ dominate, and the lemma 2.5, we obtain that there exists a finite family (α_i, β_i) , $\alpha_i \in \mathcal{H}^{SU(N,K)}(\lambda \otimes \mu^*, 0)$, $\beta_i \in \mathcal{H}^{SU(N,K)}(0, \lambda \otimes \mu^*)$, such that $\gamma_{\lambda\mu} = \langle \Omega \rangle \sum_i \alpha_i \beta_i$.

If λ and μ are distinct in $\Gamma_{N,K}$, then the two modules above are zero, hence we have $u_{\lambda\mu} = 0$.

If $\lambda = \mu$, then this two modules are generated by the duality morphisms. This gives $\sum_i \alpha_i \beta_i = \zeta d_{\lambda^*} b_{\lambda}$. We obtain $\zeta = \frac{1}{\langle \lambda \rangle}$ from the product $b_{\lambda} \gamma_{\lambda\lambda} d_{\lambda^*} = \langle \lambda \rangle \langle \Omega \rangle$. We can conclude that $u_{\lambda\lambda} = \langle \Omega \rangle$. $\square$

As already stated, using Turaev's work, modularity gives the TQFT. The universal construction of [7] could also be applied here. The normalized invariant of connected closed 3-manifolds M equipped with p_1 -structure (or 2-framing) α is defined by

$$Z(M, \alpha) = \eta \kappa^{-\sigma(\alpha)} \tau^{SU(N,K)}(M)$$

with $\kappa^3 = \Delta$. We obtain a TQFT functor (V, Z) on the cobordism category $C_{p_1}^2$ of p_1 -surfaces and (equivalence classes of) cobordisms.

Unfortunately the description of structured surfaces in [35] and [7] are not the same, however this has essentially no influence on the description of the TQFT modules.

The vectors $Z(\mathbf{D}^2 \times \mathbf{S}^1, \hat{y}_\lambda)$, $\lambda \in \Gamma_{N,K}$ form a basis for the TQFT module $V(\mathbf{S}^1 \times \mathbf{S}^1)$. Moreover this basis is orthonormal with respect to the natural hermitian form on this module.

The fusion algebra and Verlinde dimension formula. The algebra structure on the skein module of the solid torus induces the fusion algebra structure on $V(\mathbf{S}^1 \times \mathbf{S}^1)$. From modularity, we get the structure constants

$$\hat{y}_\lambda \hat{y}_\mu = \sum_{\nu} c_{\lambda\mu}^\nu \hat{y}_\nu$$

with $c_{\lambda\mu}^\nu$ equal to the rank of the module associated to a sphere, with two incoming points and one outgoing point colored respectively by λ , μ and ν . A combinatorial description for these ranks should be obtainable using [13].

For a genus g closed surface Σ_g , we can compute the dimension d_g of the TQFT module $V(\Sigma_g)$, which is equal to the invariant $Z(\Sigma_g \times \mathbf{S}^1)$. The result is given by the Verlinde formula.

$$d_g = ((N+K)^{(N-1)}N)^{g-1} \sum_{N+K > l_1 > l_2 > \dots > l_N = 0} \prod_{1 \leq i < j \leq N} \left(\frac{-1}{(s^{l_j - l_i} - s^{l_i - l_j})^2} \right)^{g-1}$$

2.4. The invariant $\tau^{PSU(N,K)}$ and the modular category $\mathcal{H}^{PSU(N,K)}$. The algebra $\mathcal{H}^{SU(N,K)}(\mathbf{D}^2 \times \mathbf{S}^1)$ is N -graded.

Set $\Gamma_{N,K}^0 = \{\lambda \in \Gamma_{N,K}, |\lambda| \equiv 0 \pmod{N}\}$.

We can see that $\Omega_0 = \sum_{\lambda \in \Gamma_{N,K}^0} \langle \lambda \rangle \hat{y}_\lambda$ satisfies the handle slide condition (an arc with a 0-graded color can slide over a component cabled with Ω_0).

Lemma 2.7. *If $d = \gcd(N, K)$ is even, and $N' = \frac{N}{d}$ $K' = \frac{K}{d}$ are both odd, then*

$$\langle U_1(\Omega_0) \rangle = 0 .$$

In all other cases one has that

$$\langle U_1(\Omega_0) \rangle \langle U_{-1}(\Omega_0) \rangle = \langle \Omega_0 \rangle = (-1)^{\frac{N(N-1)}{2}} \frac{(N+K)^{N-1}}{\prod_{j=1}^{N-1} (s^j - s^{-j})^{2(N-j)}}$$

We say that the rank-level (N, K) is *spin* if $d = \gcd(N, K)$ is even, and $N' = \frac{N}{d}$, $K' = \frac{K}{d}$ are both odd. The terminology will be justified in section 4.

We suppose now that the rank-level (N, K) is *not spin*, and that $N + K$ is invertible.

We set $\eta_0^{-2} = \langle \Omega_0 \rangle$ (we extend k if necessary), $\omega_0 = \eta_0 \Omega_0$, $\Delta = \langle U_1(\omega) \rangle$ (we can choose the sign of η_0 so that this Δ is the same as above).

Theorem 2.3. *There exists an invariant of compact oriented 3-manifolds defined on a surgery presentation by the following formula.*

$$\tau^{PSU(N,K)}(\mathbf{S}^3(L)) = \Delta^{-\sigma(L)} \langle L(\omega_0, \dots, \omega_0) \rangle$$

Remark 1. For $\gcd(N, K) = 1$ this invariant and an underlying modular category obtained from the quantum group are known (see [25]).

Remark 2. There exists a refined invariant $\tau^{SU(N,K)}(M, c)$, with c a cohomology class in $H^1(M, \mathbb{Z}/\gcd(N, K))$ such that $\tau^{PSU(N,K)}(M) = \tau^{SU(N,K)}(M, 0)$ (see section 4).

The modular category. The objects of the category $\mathbf{H}^{SU(N,K)}$ are graded by the algebraic number of points in their expansion (signed with the orientation). Let $\mathbf{H}^{PSU(N,K)}$ be the full subcategory of $\mathbf{H}^{SU(N,K)}$, whose objects are zero graded modulo N . We can show the following (we give no details).

Theorem 2.4. *The category $\mathbf{H}^{PSU(N,K)}$ is a modular category with $\Gamma_{N,K}^0$ as a representative set of isomorphism classes of simple objects.*

3. THE MODULAR CATEGORY $\tilde{\mathbf{H}}^{N,K}$ AND THE INVARIANT $\tilde{\tau}_{N,K}$

In this section, we work at rank N as before (i.e. $v = s^{-N}$), but level K will mean that, in the integral domain k ,

$$\begin{cases} s \text{ has order } 2(N + K) \text{ if } N + K \text{ is even,} \\ s \text{ has order } N + K \text{ if } N + K \text{ is odd.} \end{cases}$$

In both cases s^2 has order $N + K$; we have $v = \varepsilon s^K$, with $\varepsilon = (-1)^{N+K+1}$.

Let $d = \gcd(N, K)$, $N = dN'$, $K = dK'$. Motivated by the formulas in proposition 1.5, we would like to fix the framing parameter a in such a way that the order of the multiplicative subgroup generated by $a^N s$ and $a^K s^{-1}$ is as small as possible. Note that this order is at least d in the odd case, and $2d$ in the even case. We show that this lower bound can be realized.

To simplify the discussion we suppose in the case $N + K$ even that N' is odd; we set $d = \alpha\beta$ with $\gcd(\alpha, 2K') = \gcd(\beta, N') = 1$. Recall that $\varepsilon = (-1)^{N+K+1}$.

Lemma 3.1. *We can choose the framing parameter a so that $(a^N s)^\alpha = 1$ and $(a^K s^{-1})^\beta = \varepsilon$.*

(It may be necessary to extend the scalars.)

Proof. The problem is to find a common solution to certain polynomial equations. Here the resultant is

$$\text{Res}_a(a^{\alpha N} - s^{-\alpha}, a^{\beta K} - \varepsilon s^\beta) = (s^{-K} - \varepsilon s^N)^d$$

□

Remark 3. If $N + K$ and N' are both even, then we set $d = \alpha\beta$, with $\gcd(\alpha, K') = \gcd(\beta, N') = 1$, and in the above lemma we require that $(a^N s)^\alpha = -1$ and $(a^K s^{-1})^\beta = 1$. Following [21], we can show a level-rank duality formula, and recover this case by exchanging N and K .

In this section, and in section 4, we suppose that the framing parameter a satisfies the condition of lemma 3.1. Recall that the category $H^{(N,K,a)}$ is the quotient of the $\mathcal{C}^{N,K}$ -extended Hecke category by negligible morphisms. We define the category $\tilde{H}^{N,K}$ as follows. Objects are those of $H^{(N,K,a)}$, and the modules of morphisms are obtained from those of $H^{(N,K,a)}$ by adding as generators colored ribbon graphs in which incoming or outgoing 1-valent vertices colored with $(1^N)^{\otimes \alpha}$ or $(K)^{\otimes \beta}$ are allowed, and quotienting by the relations given by gluing an incoming $(1^N)^{\otimes \alpha}$ -colored (resp. $(K)^{\otimes \beta}$ -colored) vertex with an outgoing one. The corresponding skein functor is denoted by $\tilde{\mathcal{H}}^{N,K}$.

Exercise. a) Let $\mu \in \bar{\Gamma}_{N,K}$ be a Young diagram of the form $\mu = 1^N + \lambda$. Show that the objects μ and $1^N \otimes \lambda$ are isomorphic in the category $\tilde{H}^{N,K}$.

b) Show that the objects $(1^N)^{\otimes K}$ and $(K)^{\otimes N}$ are isomorphic in the category $\tilde{H}^{N,K}$.

c) Show that the Homflypt invariant extends to an isomorphism $\tilde{\mathcal{H}}^{N,K}(\mathbf{S}^3) \simeq k$.

Set $\dot{\Gamma}_{N,K} = \{(1^N)^{\otimes i} \otimes \lambda, 0 \leq i < \alpha \text{ and } \lambda \in \Gamma_{N,K}\}$.

The category $\tilde{H}^{N,K}$ is a ribbon category, and $\dot{\Gamma}_{N,K}$ is a finite set of dominating simple objects. We have that $\tilde{H}^{N,K}$ is a pre-modular category. This uses the involution $\star$ on the set $\dot{\Gamma}_{N,K}$ defined as follows.

For $x = (1^N)^{\otimes i} \otimes \lambda \in \dot{\Gamma}_{N,K}$, we set $x^\star = \lambda^\star \otimes (1^N)^{\otimes i'}$, where $i' \in \{0, \dots, \alpha - 1\}$ is such that the number of points in $x \otimes x^\star$ is a multiple of $N\alpha$.

We can now proceed similarly as we did in section 2.

Proposition 3.1. *Let L be a framed link in the 3-sphere, and let L' be the link obtained from L by reversing the orientation of the first component. Then for any $x \in \dot{\Gamma}_{N,K}$, one has*

$$\langle L(\hat{x}, \dots) \rangle = \langle L'(\hat{x}^*, \dots) \rangle$$

Set

$$\dot{\Omega} = \sum_{x \in \dot{\Gamma}_{N,K}} \langle x \rangle \hat{1}_x = \sum_{i=0}^{\alpha-1} \hat{g}_N^i \Omega$$

where Ω is defined as before by

$$\Omega = \sum_{\lambda \in \Gamma_{N,K}} \langle \lambda \rangle \hat{g}_\lambda$$

Proposition 3.2 (Sliding property). *The Homflypt polynomial of a link in $\mathbf{S}^3$, which has one of its components cabled with the skein element $\dot{\Omega}$, satisfies the equality in figure 2 (with Ω replaced by $\dot{\Omega}$).*

There is an action of $\mathbb{Z}/N$ on the set $\dot{\Gamma}_{N,K}$ defined as follows. The generator of $\mathbb{Z}/N$ acts by

$$(1^N)^{\otimes i} \otimes \lambda \mapsto (1^N)^{\otimes (i' + \lambda_{N-1})} \otimes ((K, \lambda) - \lambda_{N-1}^N),$$

where $i' \in \{0, \dots, \alpha - 1\}$ is congruent to $i + \lambda_{N-1}$ modulo α .

The idea is that we add to the diagram a line with K cells, and then each column which has N cells is replaced by an added copy of 1^N .

Restricting this action to the group generated by β , we obtain an action of $\mathbb{Z}/\alpha N'$. By considering the degree mod αN , we can see that this action is free. We denote by $\tilde{\Gamma}_{N,K}$ a subset of $\dot{\Gamma}_{N,K}$ which is a representative set for the orbits. We set

$$\tilde{\Omega} = \sum_{u \in \tilde{\Gamma}_{N,K}} \langle u \rangle \hat{1}_u.$$

Cabling $\dot{\Omega}$ gives the same result as cabling $N'\alpha\tilde{\Omega}$, so that $\tilde{\Omega}$ also satisfies the sliding property.

Lemma 3.2.

$$\langle U_1(\tilde{\Omega}) \rangle \langle U_{-1}(\tilde{\Omega}) \rangle = \langle \tilde{\Omega} \rangle = (-1)^{\frac{N(N-1)}{2}} \frac{d(N+K)^{N-1}}{\prod_{j=1}^{N-1} (s^j - s^{-j})^{2(N-j)}}$$

We suppose that $(N+K)$ is invertible. We set $\tilde{\eta}^{-2} = \langle \tilde{\Omega} \rangle$ (we extend k if necessary), $\tilde{\omega} = \tilde{\eta}\tilde{\Omega}$, $\delta = \langle U_1(\tilde{\omega}) \rangle$.

Theorem 3.1. *a) There exists an invariant of compact oriented 3-manifolds defined on a surgery presentation by the following formula.*

$$\tilde{\tau}_{N,K}(\mathbf{S}^3(L)) = \delta^{-\sigma(L)} \langle L(\tilde{\omega}, \dots, \tilde{\omega}) \rangle$$

b) The category $\tilde{\mathbf{H}}^{N,K}$ is a modular category with $\tilde{\Gamma}_{N,K}$ as a representative set of isomorphism classes of simple objects.

Now we give the relation between the invariant $\tau^{SU(N,K)}$ and $\tilde{\tau}_{N,K}$ in the following reduction theorem. Here $\tau^{U(1)}(M, \zeta)$ is a version of the invariant derived from linking matrices in [30] ($U(1)$ invariant) for a root of unity ζ , whose order is N' (resp. $2N'$) if N' is odd (resp. if N' is even).

Theorem 3.2 (Reduction formula). *For every manifold M , one has*

$$\tau^{SU(N,K)}(M) = \tau^{U(1)}(M, \zeta) \tilde{\tau}_{N,K}(M)$$

We will give the proof in section 5. We write this equality in the ring k where the invariant $\tau^{SU(N,K)}$ has been defined. The definition of the reduced invariant $\tilde{\tau}_{N,K}$ requires a choice of the parameters in k ; we denote by $(\tilde{s}, \tilde{v}, \tilde{a})$ this choice.

If d is even, then $\tilde{s} = s$, $\tilde{v} = v$ and $\zeta = (a^K s^{-1})^{K\beta^2}$ (the case N' even is not excluded).

If d is odd, then $\tilde{s} = -s$, $\tilde{v} = -v$ and $\zeta = ((-a)^K s^{-1})^{K\beta^2}$.

The $U(1)$ invariant in the formula above is then defined by

$$\tau^{U(1)}(\mathbf{S}^3(L), \zeta) = \left(\frac{\Delta}{\delta}\right)^{-\sigma(L)} \left(\frac{\eta}{\tilde{\eta}}\right)^m \sum_{j \in (\mathbb{Z}/N')^m} \zeta^{t_{jB_L j}}.$$

4. REFINED INVARIANTS

4.1. Spin structures modulo an even integer. In [29], H. Murakami stated a decomposition formula for the invariant $\tau^{SU(N,K)}$ using some spin type structures (see remark 2.7 in his paper). He observed that for $N = 2$ these are spin structures, and the corresponding refinements were studied in [19] and [5]. For $N > 2$ he only gave a combinatorial description of the structures, and asked for a topological interpretation. We recall here the topological definition for these structures which we gave in [6].

Suppose d is an even integer. Then there exists, up to homotopy, a unique non trivial map $g : BSO \rightarrow K(\mathbb{Z}/d, 2)$. Define the fibration

$$\pi_d : BSpin(\mathbb{Z}/d) \rightarrow BSO$$

to be the pull-back, using g , of the path fibration over $K(\mathbb{Z}/d, 2)$. The space $BSpin(\mathbb{Z}/d)$ is a classifying space for the non trivial central extension of the Lie group SO by $\mathbb{Z}/d$, which we denote by $Spin(\mathbb{Z}/d)$.

Now we can use the fibration π_d to define structures (see [St]). Let $\gamma_{Spin(\mathbb{Z}/d)} = \pi_d^*(\gamma_{SO})$ be the pull-back of the canonical vector bundle over BSO .

Definition 4.1. A spin structure with mod d coefficients on a manifold M is an homotopy class of fiber maps from the stable tangent bundle τ_M to $\gamma_{Spin(\mathbb{Z}/d)}$.

If non empty, the set of these structures, denoted by $Spin(M; \mathbb{Z}/d)$, is affinely isomorphic to $H^1(M; \mathbb{Z}/d)$, by obstruction theory. Moreover the obstruction for existence is a class $w_2(M; \mathbb{Z}/d) \in H^2(M; \mathbb{Z}/d)$, which is the image of the Stiefel-Whitney class $w_2(M)$ by the homomorphism induced by the inclusion of coefficients $\mathbb{Z}/2 \hookrightarrow \mathbb{Z}/d$. As the Stiefel-Whitney class $w_2(M)$ is zero for every compact oriented 3-manifold, this shows that spin structures modulo d exist on every 3-manifold M . The following theorem gives a combinatorial description for these structures. Recall that a surgered manifold $M = \mathbf{S}^3(L)$ is the boundary of a 4-manifold W_L called the trace of the surgery. To each $\sigma \in Spin(M; \mathbb{Z}/d)$ is associated a relative obstruction $w_2(\sigma; \mathbb{Z}/d)$ in $H^2(W_L, M; \mathbb{Z}/d)$. The group $H^2(W_L, M; \mathbb{Z}/d)$ is free of rank $m = \sharp L$. Taking the coordinates of the relative obstruction we get a map $\psi_L : Spin(M; \mathbb{Z}/d) \rightarrow (\mathbb{Z}/d)^m$.

Theorem 4.1. *The map ψ_L is injective, and its image is the set of those $(c_1, \dots, c_m)$ which are solutions of the following $(\mathbb{Z}/d)$ -characteristic equation*

$$B_L \begin{pmatrix} c_1 \\ \vdots \\ c_m \end{pmatrix} = \frac{d}{2} \begin{pmatrix} b_{11} \\ \vdots \\ b_{mm} \end{pmatrix} \pmod{d}.$$

Here the b_{ii} are the diagonal values of the linking matrix B_L .

Proof. First we compute the absolute obstruction $w_2(W_L; \mathbb{Z}/d) = \xi_*(w_2(W_L))$, where ξ_* is induced by the morphism of coefficients $\xi : \mathbb{Z}/2 \hookrightarrow \mathbb{Z}/d$. If x is an integral 2-cycle in W_L with self-intersection $x.x$ and $[x]_\nu$ denotes its homology class modulo an integer ν , $w_2(W_L) \in H^2(W_L; \mathbb{Z}/2)$ is determined by the equation

$$\forall x < w_2(W_L), [x]_2 > = x.x \pmod{2}.$$

Hence $w_2(W_L; \mathbb{Z}/d) \in H^2(W_L; \mathbb{Z}/d)$ is determined by

$$\forall x < w_2(W_L; \mathbb{Z}/d), [x]_d > = \xi(x.x) = \frac{d}{2}x.x \pmod{d}.$$

Now by functoriality, the relative obstruction lives in the inverse image of the absolute one under the map induced by inclusion $H^2(W_L, M; \mathbb{Z}/d) \rightarrow H^2(W_L; \mathbb{Z}/d)$. Using the affine structure over $H^1(M; \mathbb{Z}/d)$, we obtain an affine bijection between $Spin(M; \mathbb{Z}/d)$ and this inverse image. Whence we have the lemma by writing the equation above using coordinates. $\square$

There is a formula for the bijection $\psi_{L,L'}$ corresponding to a Kirby move. Using the $\mathbb{Z}/d$ -characteristic equation we see that the coefficient for a trivial component with framing ± 1 is $d/2$. For the usual positive Fenn-Rourke move, the formula is

$$\psi_{L,L'}(c_1, \dots, c_{m-1}, d/2) = (c_1, \dots, c_{m-1}, c'_m)$$

$$\text{with } c'_m = d/2 - \sum_i b'_{im} c_i.$$

Here b'_{im} is the (i, m) -indexed coefficient of the matrix $B_{L'}$.

4.2. Spin refinements. We consider here the reduced theory of section 3 in the spin case. Recall that this means

$$\begin{cases} d = \gcd(N, K) \text{ is even,} \\ N' = \frac{N}{d} \text{ and } K' = \frac{K}{d} \text{ are odd.} \end{cases}$$

We decompose the skein element $\tilde{\omega} = \sum_{\nu} \tilde{\omega}_{\nu}$ according to the $\mathbb{Z}/d$ -grading of the algebra $\tilde{\mathcal{H}}^{N,K}(\mathbf{D}^2 \times \mathbf{S}^1) = \bigoplus_{\nu} \tilde{\mathcal{H}}_{\nu}^{N,K}(\mathbf{D}^2 \times \mathbf{S}^1)$.

Theorem 4.2. *Provided $c = (c_1, \dots, c_m)$ satisfies the modulo d characteristic condition, the formula*

$$\tilde{\tau}_{N,K}^{\text{spin}}(M, \sigma) = \delta^{-\sigma(L)} \langle L(\tilde{\omega}_{c_1}, \dots, \tilde{\omega}_{c_m}) \rangle$$

defines an invariant of the surgered manifold $M = \mathbf{S}^3(L)$ equipped with the modulo d spin structure $\sigma = \psi_L^{-1}(c_1, \dots, c_m)$.

Moreover,

$$\forall M \quad \tilde{\tau}_{N,K}(M) = \sum_{\sigma \in Spin(M; \mathbb{Z}/d)} \tilde{\tau}_{N,K}^{\text{spin}}(M, \sigma).$$

Proof. The following graded version of the sliding property can be derived from the proof of proposition 2.3.

Lemma 4.1 (Graded sliding property). *The Homflypt polynomial of a link in $\mathbf{S}^3$, which has one of its components cabled with the skein element $\tilde{\omega}_{\nu}$, satisfies the equality in figure 4.*




FIGURE 4. Graded sliding property

Using this sliding property, we get

$$\forall \nu \forall x_\nu \in \tilde{\mathcal{H}}_\nu^{N,K}(\mathbf{D}^2 \times \mathbf{S}^1) \langle H_1(x_\nu, \tilde{\omega}_{d/2-\nu}) \rangle = \langle U_0(x_\nu) \rangle \langle U_1(\tilde{\omega}_{d/2}) \rangle$$

This shows that the spin handle slide condition is satisfied.

We can deduce from the lemma 4.3 that $\langle U_\epsilon(\tilde{\omega}_{d/2}) \rangle = \langle U_\epsilon(\tilde{\omega}) \rangle$ is invertible, and hence we have that the invariant $\tilde{\tau}_{N,K}^{\text{spin}}$ is well defined.

Now we show the decomposition formula. We can write

$$\langle L(\tilde{\omega}, \dots, \tilde{\omega}) \rangle = \sum_c \langle L(\tilde{\omega}_{c_1}, \dots, \tilde{\omega}_{c_m}) \rangle.$$

The result is contained in the lemma 4.2. $\square$

Lemma 4.2. *If c does not satisfy the modulo d characteristic condition, then*

$$\langle L(\tilde{\omega}_{c_1}, \dots, \tilde{\omega}_{c_m}) \rangle = 0.$$

Proof. Let $L = (L_1, \dots, L_m)$. Up to a permutation of the components, we have to show that $\langle L(\tilde{\omega}_{c_1}, \dots, \tilde{\omega}_{c_m}) \rangle = 0$, if $\sum_{j=1}^m b_{1j}c_j \not\equiv \frac{d}{2} \pmod{d}$. The proof is in three steps.

If L_1 is a 1-framed unknot, then, using the sliding property, we get the result from the lemma 4.3.


If L_1 is an unknot with any framing, note that (up to an invertible factor) we can add to the link a trivial ± 1 -framed U_ϵ cabled with $\omega_{d/2}$; we then use the sliding property to move this unknot around L_1 , and this adds ϵ to the framing. This allows us to reduce the problem to the preceding case.

In the general case, the component L_1 can be unknotted by changing some crossings, and inserting a ± 1 -framed unknot cabled with $\omega_{d/2}$ around the crossing, in such a way that its linking number with L_1 is zero. By the sliding property, this amounts to multiplying by an invertible element, whence we have the result by using the above. $\square$

Lemma 4.3. *For $\epsilon = \pm 1$,*

$$\langle U_\epsilon(\tilde{\omega}_\nu) \rangle = 0 \text{ if } \nu \neq \frac{d}{2}.$$

Proof. We compute $\langle U_1(\tilde{\omega}_\nu) \rangle \langle U_{-1}(\tilde{\omega}_\nu) \rangle$. The graded sliding property shows that this is equal to the Homflypt invariant of the following colored link.




Using the vanishing lemma 4.4 below, whose proof is adapted from the one given in 2.5, we obtain

$$\langle U_1(\tilde{\omega}_\nu) \rangle \langle U_{-1}(\tilde{\omega}_\nu) \rangle = \tilde{\eta} \sum_{i=0}^{\alpha-1} (a^N s)^{2\nu i} \sum_{j=0}^{\beta-1} (-1)^j (a^K s^{-1})^{2\nu j}.$$

If $\langle U_1(\tilde{\omega}_\nu) \rangle \langle U_{-1}(\tilde{\omega}_\nu) \rangle$ is not zero, then we have $\nu \equiv 0 \pmod{\alpha}$, and $\nu \equiv \frac{\beta}{2} \pmod{\beta}$. $\square$

Lemma 4.4. *If λ is a Young diagram in $\Gamma_{N,K} \setminus \{K^j, 0 \leq j < N\}$, then, for any $\nu \in \mathbb{Z}/d$, the following morphism is zero in $\tilde{H}^{N,K}$.*



4.3. Cohomological refinements. If the rank-level is not spin, we can proceed similarly. This time we have that $\langle U_1(\tilde{\omega}_\nu) \rangle = 0$, unless $\nu = 0 \pmod{d}$.

Theorem 4.3. *Provided $c = (c_1, \dots, c_m)$ is in the kernel of the linking matrix modulo d , the formula*

$$\tilde{\tau}_{N,K}^{\text{coho}}(M, \sigma) = \delta^{-\sigma(L)} \langle L(\tilde{\omega}_{c_1}, \dots, \tilde{\omega}_{c_m}) \rangle$$

defines an invariant of the surgered manifold $M = \mathbf{S}^3(L)$ equipped with the cohomological $\sigma \in H^1(M, \mathbb{Z}/d)$ corresponding to c .

Moreover,

$$\forall M \quad \tilde{\tau}_{N,K}(M) = \sum_{\sigma \in H^1(M; \mathbb{Z}/d)} \tilde{\tau}_{N,K}^{\text{coho}}(M, \sigma)$$

5. PROOF OF THE REDUCTION FORMULA

We show the reduction formula (theorem 3.2) in the spin case. We proceed as follows. We use the modulo β grading to construct, as we did for $\tilde{\tau}_{N,K}$, a spin refinement of the invariant $\tau^{SU(N,K)}$ satisfying the decomposition formula

$$\tau^{SU(N,K)}(M) = \sum_{\sigma \in \text{Spin}(M, \mathbb{Z}/\beta)} \tau^{SU(N,K)}(M, \sigma)$$

The same can be done with the reduced invariant $\tilde{\tau}_{N,K}$. Note that here we consider the modulo β grading, so that the decomposition formula below is not that of the preceding section if $\alpha \neq 1$.

$$\tilde{\tau}_{N,K}(M) = \sum_{\sigma \in \text{Spin}(M, \mathbb{Z}/\beta)} \tilde{\tau}_{N,K}(M, \sigma)$$

We will prove the reduction theorem for the spin invariants. We need to specify some notation. We will use $\langle \rangle$ for the Homflypt invariant evaluated at a , and $\langle \rangle^\sim$ for the Homflypt invariant evaluated at $\tilde{a}$. Note that the framing parameter does not appear in the coefficients of $\dot{\Omega}$, so that we can use them for cabling and then evaluate the Homflypt invariant at a or $\tilde{a}$.

We decompose the skein element $\dot{\Omega}$ according to the $N\alpha$ -grading (resp. the β -grading).

$$\dot{\Omega} = \sum_{\xi=0}^{\alpha N-1} \dot{\Omega}^{(\xi)} \quad \left(\text{resp. } \dot{\Omega} = \sum_{\nu=0}^{\beta-1} \dot{\Omega}_\nu \right)$$

Formulas for the modulo β spin invariants are then

$$\tau^{SU(N,K)}(\mathbf{S}^3(L), \sigma_c) = \Delta^{-\sigma(L)} (\alpha^{-1} \eta)^m \langle L(\dot{\Omega}_{c_1}, \dots, \dot{\Omega}_{c_m}) \rangle$$

$$\tilde{\tau}_{N,K}(\mathbf{S}^3(L), \sigma_c) = \delta^{-\sigma(L)} (\alpha^{-1} N'^{-1} \tilde{\eta})^m \langle L(\dot{\Omega}_{c_1}, \dots, \dot{\Omega}_{c_m}) \rangle^\sim$$

Here $c = (c_1, \dots, c_m)$ satisfies the modulo β characteristic condition, and σ_c is the modulo β spin structure corresponding to c . We can choose L so that the linking matrix is even. (This can be considered to be a consequence of the nullity of the cobordism group Ω_3^{spin} .) In this case c is *modulo β characteristic* if and only if c is in the kernel of the linking matrix modulo β .

Lemma 5.1. *We can find $\xi_1, \dots, \xi_m \in \mathbb{Z}/\alpha N$ such that $\xi_i \equiv c_i \pmod{\beta}$, for $i = 1, \dots, m$ and $\xi = (\xi_1, \dots, \xi_m)$ represents an element in the kernel of the linking matrix modulo αN .*

Proof. Denote by B the linking matrix. The Bockstein operator

$$\text{Ker}(B \otimes \mathbb{Z}/\beta) \rightarrow \text{coker}(B \otimes \mathbb{Z}/\alpha^2 N')$$

associated with the exact sequence

$$0 \rightarrow \mathbb{Z}/\alpha^2 N' \rightarrow \mathbb{Z}/\alpha N \rightarrow \mathbb{Z}/\beta \rightarrow 0$$

is zero since $\gcd(\beta, \alpha^2 N') = 1$. $\square$

In the formula for $\tau^{SU(N,K)}(\mathbf{S}^3(L), \sigma_c)$ and $\tilde{\tau}_{N,K}(\mathbf{S}^3(L), \sigma_c)$ we can replace $\dot{\Omega}_{c_i}$ by $\sum_{j=0}^{\alpha^2 N' - 1} \hat{y}_{K\beta}^j \dot{\Omega}^{(\xi_i)}$ (with ξ fixed as in the lemma above). Using the braiding and framing coefficients for $y_{K\beta}$, we obtain the formula

$$\begin{aligned} \langle L(\dot{\Omega}_{c_1}, \dots, \dot{\Omega}_{c_m}) \rangle &= \sum_{j \in (\mathbb{Z}/\alpha^2 N')^m} \zeta^{t j B_L j} \langle L(\dot{\Omega}^{(\xi_1)}, \dots, \dot{\Omega}^{(\xi_m)}) \rangle \\ \langle L(\dot{\Omega}_{c_1}, \dots, \dot{\Omega}_{c_m}) \rangle &= \alpha^{2m} \sum_{j \in (\mathbb{Z}/N')^m} \zeta^{t j B_L j} \langle L(\dot{\Omega}^{(\xi_1)}, \dots, \dot{\Omega}^{(\xi_m)}) \rangle \end{aligned}$$

We also have

$$\langle L(\dot{\Omega}_{c_1}, \dots, \dot{\Omega}_{c_m}) \rangle^\sim = (\alpha^2 N')^m \langle L(\dot{\Omega}^{(\xi_1)}, \dots, \dot{\Omega}^{(\xi_m)}) \rangle^\sim$$

For any framed link K , we have $\langle K \rangle = \left(\frac{a}{\delta}\right)^{K.K} \langle K \rangle^\sim$. Using that $\left(\frac{a}{\delta}\right)^{\alpha N} = 1$, we see that

$$\langle L(\dot{\Omega}^{(\xi_1)}, \dots, \dot{\Omega}^{(\xi_m)}) \rangle = \langle L(\dot{\Omega}_{c_1}, \dots, \dot{\Omega}_{c_m}) \rangle^\sim$$

We deduce the required formula with the following normalization of the $U(1)$ invariant at the N' -th root of unity ζ .

$$\tau^{U(1)}(\mathbf{S}^3(L), \zeta) = \left(\frac{\Delta}{\delta}\right)^{-\sigma(L)} \left(\frac{\eta}{\tilde{\eta}}\right)^m \sum_{j \in (\mathbb{Z}/N')^m} \zeta^{t j B_L j}$$

Note that $g = \frac{\Delta}{\delta} \frac{\eta}{\tilde{\eta}}$ is a Gauss sum, whose square modulus is $g\bar{g} = \left(\frac{\eta}{\tilde{\eta}}\right)^2 = N'$.

The other cases are obtained similarly. When $N + K$ is odd it is useful to note first that in the defining expression for $\tau^{SU(N,K)}$ we can evaluate the Homflypt invariant at $(-a, -s, -v)$ as well as at (a, s, v) .

REFERENCES

1. A. K. Aiston, *Skein theoretic idempotents of Hecke algebras and quantum group invariants*, Thesis, University of Liverpool, 1996.
2. A. K. Aiston, *A skein theoretic proof of the hook formula for quantum dimension*, (preprint 1997).
3. A. K. Aiston, H.R. Morton, *Idempotents of Hecke algebras of type A*, (preprint 1997).
4. H. H. Andersen, *Tensor products of quantized tilting modules*, Comm. Math. Phys. 149 (1991), 149–159.
5. C. Blanchet, *Invariants on three-manifolds with spin structure*, Comm. Math. Helv. 67(1992), 406–427.
6. C. Blanchet, *Refined quantum invariants for three-manifolds with structure*, in Knot Theory, Banach Center Pub. Vol 42, to appear.
7. C. Blanchet, N. Habegger, G. Masbaum and P. Vogel, *Topological Quantum Field Theories derived from the Kauffman bracket*, Topology 34 No 4 (1995), 883–927.
8. N. Bourbaki, *Groupes et algèbres de Lie IV, V, VI*, Masson, Paris (1982).
9. A. Bruguières, *Tresses et structure entière sur la catégorie des représentations de $Sl_{N,q}$* (preprint 1998).
10. G. Dipper, G. James, *Block idempotents of the Hecke algebra of the general linear group*, Proc. London Math. Soc. 54 (1987) 57–82.
11. J. Erlijman, *New subfactors from braid group representations*, (preprint 1997).
12. M. Geck, *Representations of Hecke algebras at roots of unity*, Séminaire Bourbaki No 836, Nov 1997.
13. F. M. Goodman, H. Wenzl, *Littlewood-Richardson Coefficients for Hecke algebras at Roots of Unity*, Advances in Math. 82 (1990), 244–265.
14. A. Gyoja, *A q -analogue of Young symmetrizer*, Osaka J. Math. 23 (1986), 841–852.
15. P. Freyd, D. Yetter, J. Hoste, W. B. R. Lickorish, K. Millet and A. Ocneanu, *A new polynomial invariant of knots and links*, Bull. AMS 12 (1985), 239–246.
16. V. Jones, *Hecke algebra representations of braid groups and link polynomials*, Annals of math. 126 (1987), 335–338.
17. D. Kazhdan, H. Wenzl, *Reconstructing monoidal categories*, Adv. in Soviet Math. 16 n. 2 (1993), 111–136.
18. R. Kirby, *A calculus of framed links in S^3* , Invent. Math. 45 (1978), 35–56.
19. R. Kirby, P. Melvin, *The 3-manifold invariants of Witten and Reshetikhin-Turaev for $sl(2, \mathbb{C})$* , Inv. Math. 105, 473–545 (1991).
20. T. Kohno, T. Takata, *Symmetry for Witten’s 3-manifold invariants for $sl(n, \mathbb{C})$* , Journal of Knot Theory and its Ram. 2 (1993), 149–169.
21. T. Kohno, T. Takata, *Level-rank duality for Witten’s 3-manifold invariant*, Adv. Studies Pure Math. 24, Progress in Algebraic Combinatorics, 243–264.
22. W. B. R. Lickorish, *Sampling the $SU(N)$ invariants of three-manifolds*, J. of Knot Theory and Ram. (1997).
23. W. B. R. Lickorish, *Skeins, $SU(N)$ three-manifolds invariants and TQFT*, (preprint 1997).
24. I. G. Macdonald, *Symmetric functions and Hall polynomials*, Oxford Math. Monographs (1979).

25. G. Masbaum, H. Wenzl, *Integral modular categories and integrality of quantum invariants at roots of unity of prime order*, (preprint 1997).
26. H.R.Morton, *Invariants of links and 3-manifolds from skein theory and from quantum groups*, in M. Bozhyk, editor, *Topics in knot theory*, N.A.T.O. A.S.I. series C 399, Kluwer (1993), 107–156.
27. H. R. Morton and A. K. Aiston, *Young diagrams, the Homfly skein of the annulus and unitary invariants*, (preprint 1996).
28. H. R. Morton and P. Traczyk, *Knots and algebras*, Contribuciones Matematicas en homenaje al professor D. Antonio Plans Sanz de Bremond, E. Martin-Peinador and A. Rodez editors, University of Saragoza (1990), 201–220.
29. H. Murakami, *Quantum invariants for 3-manifolds*, Proc. of the 3rd Korea-Japan School of knots and links.
30. H. Murakami, T. Ohtsuki and M. Okada, *Invariants of three-manifolds derived from linking matrices of framed links*, Osaka J. Math. 29 (1992), 545–572.
31. J. Przytycki and P. Traczyk, *Invariants of links of Conway type*, Kobe J. Math. 4 (1987), 115–139.
32. A. Ram, H. Wenzl, *Matrix units for centralizer algebras*, J. of Alg. 145 (1992), 378–395.
33. N. Reshetikhin and V. Turaev, *Invariants of 3-manifolds via link polynomials and quantum groups*, Invent. Math. 103 (1991), 547–598.
34. V. G. Turaev, *Operator invariants of tangles, and R-matrices*, Math. USSR Izv. Vol. 35 (1990), No 2, 411–443.
35. V. G. Turaev, *Quantum invariants of knots and 3-manifolds*, De Gruyter Studies in Math. 18 (1994).
36. V. Turaev, H. Wenzl, *Quantum invariants of 3-manifolds associated with classical simple Lie algebras*, Int. J. of Math. Vol 4, No 2 (1993) 323–358.
37. V. Turaev, H. Wenzl, *Semisimple and modular categories from link invariants*, Math. Ann. 309 (1997), 411–461.
38. P. Vogel, *Les invariants récents des variétés de dimension 3*, Séminaire Bourbaki 799 (1995).
39. H. Wenzl, *Hecke algebras of type A_n and subfactors*, Invent. Math. 92 (1988), 349–383.
40. H. Wenzl, *Braids and invariants of 3-manifolds*, Invent. Math. 114 (1993), 235–275.
41. Y. Yokota, *Skeins and quantum $SU(N)$ invariants of 3-manifolds*, Math. Ann. 307 (1997), 109–138.

UNIVERSIT DE NANTES-UMR 6629, 2 RUE DE LA HOUSSINIRE, BP92208,
 44322 NANTES CEDEX 3, FRANCE
E-mail address: blanchet@math.univ-nantes.fr